



» NON-ALCOHOL RAPID SURFACE DISINFECTANTS «

# ACTOANID® WIPES

Ready-to-Use Disinfection Wipes for Alcohol-Sensitive Surfaces

- ✓ A combination of modern active ingredients: PHMB and QAC
- ✓ Suitable to use on surfaces sensitive to alcohol
- ✓ Alcohol, aldehyde and phenol-free
- ✓ High material compatibility including incubators and acrylic glasses
- ✓ Effective in 60 sec. against bactericidal, fungicidal and limited virucidal
- ✓ Ready-to-use



**Actoanid® Wipes** is used for cleaning and disinfection of the surfaces of alcohol-sensitive medical devices which are not suitable for contact with liquid. Polyhexanide and quaternary ammonium combination based. It is a product suitable for cleaning large areas and it is in the form of wipes containing nonwovens (viscose/polyester). It does not contain aldehyde and phenol derivatives.



German Company



**ACTO®** Pharma  
Cosmetic  
Disinfection  
Wound Care  
Food Supplement

Active Chemical Technology Organisation

# ACTOANID® WIPES



## Ready-to-Use Disinfection Wipes for Alcohol-Sensitive Surfaces

### Field of Application:

It is used for cleaning and disinfection of medical devices with a non-invasive, regular and smooth surface. Suitable for safe and effective disinfection of medical devices such as incubators, ultrasound probes (excluding vaginal probes), external surfaces of hemodialysis devices, dental units (excluding drills and hoses) that need to be disinfected frequently.

### Usage:


Remove the blood and other body fluids on the surface with an absorbent material before the procedure. Wipe the surface of the medical device with **Actoanid® Wipes** to use 3-4 wipes per square meter.

Make sure that the wipes reach the entire surface of the device and there is no dry place left. Wait for the duration of the effect. Do not dry after use. The packaging should be tightly closed after each use. Used wipes should not be reused.


### Chemical Composition:

Content of 100 g Impregnation Solution:  
0.25 g Poly (Hexamethylene Biguanide) Hydrochloride, (PHMB), 0.25 g Quaternary Ammonium Compounds (QAC) and excipients.


### Instructions For Use:




Wear gloves




Open the flap lid




Open the foil




Wear gloves




Remove the cap




Remove the foil




Open the hinged lid




Take a wipe



Close the foil



Close the lid tightly




Put the towel through the hinged lid



Fasten the cap



Tear off the towel



Fasten the hinged lid

Packaging		
Unit	Pieces in a Box	Article-No.
80 Wipes - 15x20 cm	18 (Packaging Bag)	05.1210.80
100 Wipes - 15x24 cm	6 (PE Container)	05.1210.100

Microbiological Activity	Test Method	Concentration	Effect Time
Bactericidal	EN 16615	Ready-to-use	1 min.
Fungicidal ( <i>Aspergillus brasiliensis</i> , <i>Candida albicans</i> )	EN 16615	Ready-to-use	1 min.
Limited Virucidal ( <i>Vaccinia</i> )	EN 14476	Ready-to-use	1 min.

